

TABLE 1.—*Causes of Neurogenic Diabetes Insipidus in the Neonatal Period*

| |
|-----------------------------|
| Intracranial defects |
| Holoprosencephaly |
| Septo-optic dysplasia |
| Infections |
| Listerial sepsis |
| Meningitis—streptococcal |
| Meningococcal septicemia |
| Intracranial tumors |
| Trauma |
| Intraventricular hemorrhage |
| Brain death |

of 121 nmol per liter (4.4 μ g per dl [normal, 110 to 520 nmol per liter]), thyroid-stimulating hormone level of 3.9 mU per liter, an unenhanced brain computed tomographic scan showing no intracranial defects, normal optic nerves and septum pellucidum, and normal plain films of the skull and long bones.

The infant had complete resolution of vomiting and lethargy, and the findings of a physical examination on discharge were completely within normal limits. She was feeding well on formula and was sent home on a regimen of intranasal desmopressin. At 12 weeks of age she was noted to have poor response to visual stimuli. An ophthalmologic examination revealed nystagmus and hypoplastic optic nerves—approximately 10% normal size. The diagnosis of de Morsier's syndrome (septo-optic dysplasia) was made.

Discussion

Neurogenic diabetes insipidus of neonates usually presents in a patient with known predisposing risk factors (Table 1). As such, the neonate is typically being treated or followed for the preexisting conditions, and diabetes insipidus is discovered because of abnormal laboratory values or urine output. The patient we describe presented with diabetes insipidus at 21 days of age with no known predisposing conditions, and an initial workup proved uninformative regarding a cause. Of interest is that the baby reportedly preferred water over formula since birth, although previous physical examinations revealed no signs of dehydration. Additionally, eye movements and gross visual responses were noted to be normal in the newborn nursery. A further evaluation at 3 months of age revealed diminished visual responses and findings consistent with de Morsier's syndrome (septo-optic dysplasia).

The de Morsier's syndrome is caused by congenital midline brain malformations thought to occur at 6 to 8 weeks of gestational age and involves cerebral structures, optic nerves, pituitary gland, and the hypothalamus to a variable degree. It usually affects the offspring of young primigravid mothers (which was the case in this patient). Unlike the patient described here, infants with de Morsier's syndrome usually present with ophthalmologic abnormalities as the chief parental concern.⁵ The endocrine abnormalities are found subsequently and can be multiple or isolated, although growth hormone deficiency is the most consistent abnormality. Approximately 40% of patients will have an absence of the septum pellucidum, although its presence or absence does not correlate with endocrine abnormalities.⁶ Again, the patient we described had a normal septum pellucidum.

This case exemplifies the need for continued follow-up to

determine the cause of central diabetes insipidus in an infant once the diagnosis has been made. Greger and co-workers reported that in a series of 58 children, the cause of diabetes insipidus could eventually be found in more than 90% of patients.⁷ An ongoing evaluation of patients with diabetes insipidus in whom no obvious cause is evident should include a computed tomographic scan of the brain at intervals, repeated evaluations of the hypothalamic-pituitary axis, and astute physical examinations.

REFERENCES

1. Jenkins HR, Hughes IA, Gray OP: Cranial diabetes insipidus in early infancy. *Arch Dis Child* 1988; 63:434-435
2. Kappy MS, Sonderer E: Sublingual administration of desmopressin—Effectiveness of an infant with holoprosencephaly and central diabetes insipidus. *Am J Dis Child* 1987; 141:84-85
3. Adams JM, Kenny JD, Rudolph AJ: Central diabetes insipidus following intraventricular hemorrhage. *J Pediatr* 1976; 88:292-294
4. Fenton LJ, Kleinman LI: Transient diabetes insipidus in a newborn infant. *J Pediatr* 1974; 85:79-81
5. Izenberg N, Rosenblum M, Parks JS: The endocrine spectrum of septo-optic dysplasia. *Clin Pediatr* 1984; 23:632-636
6. Stanhope R, Preece MA, Brook CGD: Hypoplastic optic nerves and pituitary dysfunction. *Arch Dis Child* 1984; 59:111-114
7. Greger NG, Kirkland RT, Clayton GW, et al: Central diabetes insipidus—22 years' experience. *Am J Dis Child* 1986; 140:551-554

Occult Retroperitoneal Carcinoid Tumor With Flushing and Solitary Lung Metastasis

JEFFREY A. JACKSON, MD
Temple, Texas

HOPE D. SHIPMAN, MD
Houston

FLUSHING is the hallmark symptom of the carcinoid syndrome, occurring in 63% to 96% of affected patients.¹⁻⁴ By the time of overt symptoms, patients with carcinoid syndrome usually have liver metastases and pronounced elevations in urinary 5-hydroxyindoleacetic acid (5-HIAA), and the diagnosis is straightforward. We describe the diagnostic challenge of a patient with flushing who had initially normal to minimally elevated urinary 5-HIAA levels and a small pulmonary nodule and later was shown to have an occult metastatic carcinoid tumor of apparent retroperitoneal origin. His dramatic response to the long-acting somatostatin analogue, compound SMS-201-995 (octreotide [Sandostatin, Sandoz Research Institute, East Hanover, NJ]), with relief of subsequent intractable abdominal pain is also reported.

Report of a Case

The patient, a 58-year-old man, had mild facial flushing and intermittently loose stools in April 1985. That fall, his internist measured 24-hour urinary 5-HIAA concentrations on three occasions, with values of 40, 43, and 60 μ mol per day (normal 10 to 52), and abdominal computed tomography

(Jackson JA, Shipman HD: Occult retroperitoneal carcinoid tumor with flushing and solitary lung metastasis. *West J Med* 1989 Oct; 151:461-463)

From the Department of Medicine, Division of Endocrinology, Scott and White Clinic, Scott and White Memorial Hospital, Scott, Sherwood and Brindley Foundation, Texas A&M University College of Medicine, Temple. Dr Shipman is currently with the McGreggor Medical Association, Houston.

Reprint requests to Jeffrey A. Jackson, MD, Division of Endocrinology, Department of Medicine, Scott and White Clinic, 2401 S 31st St, Temple, TX 76708.

ABBREVIATIONS USED IN TEXT

CT = computed tomography

5-HIAA = 5-hydroxyindoleacetic acid

(CT) showed mild retrocrural and periaortic adenopathy with no evidence of liver metastasis. His initial chest x-ray film was normal. A urinary histamine level was slightly elevated at 75 μg per 24 hours (normal 17 to 68), but a bone marrow aspiration biopsy specimen did not show mastocytosis. The blood serotonin level (SmithKline Bio-Science Laboratories, Van Nuys, Calif) was 2.51 μmol per liter (normal 0.28 to 1.14), and a plasma calcitonin level was 6 ng per liter (normal < 19). Vasoactive intestinal polypeptide and substance P levels (A. I. Vinik, MD, University of Michigan Medical School, Ann Arbor) were undetectable—less than 10 ng per liter and less than 25 ng per liter, respectively. A follow-up abdominal CT scan six months later was unchanged; a chest CT scan with 0.5-cm cuts showed a 6-mm right inferior lung field nodule. Serotonin levels gradually rose, yet urinary 5-HIAA concentrations remained normal. In early 1987, a chest x-ray film showed that the nodule had increased in size to 8 to 9 mm with unchanged findings on an abdominal CT scan. No evidence suggested that the minor abdominal lymphadenopathy (nodes ≤ 1 cm), which had remained unchanged over a two-year period, represented carcinoid metastases. The patient finally consented to wedge resection of the right middle lobe. A localized carcinoid nodule was removed, but postoperatively his flushing persisted.

Intestinal pseudo-obstruction with abdominal pain occurred three months later, and for the first time small liver metastases were noted by CT and magnetic resonance imaging. Angiography showed extensive mesenteric encasement and splenic artery occlusion consistent with an unresectable infiltrative mass lesion deep at the root of the mesentery in the region of the pancreatic head (not visualized by CT or magnetic resonance imaging). Large venous varicosities draining into the caval system were noted in the retroperitoneal area.

Intractable abdominal pain ensued with nonobstructive nausea, vomiting, and weight loss, and the 24-hour urinary concentration of 5-HIAA increased to 345 μmol . Octreotide therapy was begun in August 1987, and the dosage was gradually increased to 200 μg subcutaneously twice a day. Within two months, his abdominal pain decreased considerably, with a reduction in flushing and diarrhea to once or twice a month. As of November 1988, he had gained 8 kg (18 lb) and was free of flushing and abdominal pain with the discontinuation of narcotic analgesia ten months before. Urinary 5-HIAA concentrations fell to 40 μmol per day. A follow-up CT scan showed no growth of liver metastasis; no additional metastases were noted. The only complications of octreotide therapy were mild steatorrhea—16.8 grams per 24 hours (normal < 7.0)—and gallstones noted by sonography after nine months of treatment; both have remained asymptomatic. Interestingly, dysosmia, which was noted by the patient early in his course, diminished markedly during octreotide therapy.

Discussion

The differential diagnosis of flushing is diverse,^{5,6} and the clinical evaluation can be challenging. Provocative testing with ethanol, epinephrine, or pentagastrin^{2,7-9} can be done to

precipitate flushing in doubtful circumstances, but the results are not specific⁵ and the test may induce dangerous tachycardia or hypotension. When used in combination with measuring specific markers such as tachykinins, provocative testing may be diagnostic.⁹

If the clinical suspicion warrants, a search for other hormonal markers should be undertaken in a patient with flushing and normal 5-HIAA levels. Blood serotonin is an excellent marker of carcinoid hypersecretion,^{6,8,10-12} as shown in this case. Urinary serotonin^{10,12,13} and histamine,^{2,10} circulating levels of substance P,^{9,11,12} chromogranin,¹⁴ neurotensin,¹² and other tachykinins,⁹ and platelet serotonin levels^{12,13,15} may also have value as carcinoid markers. Selective tumoral production of 5-hydroxytryptophan, particularly by foregut carcinoids, has been reported,^{2,16} but levels of urinary 5-HIAA and serotonin are usually abnormal in these patients^{13,16} because of a peripheral conversion of 5-hydroxytryptophan to serotonin and 5-HIAA. Less than 1% of carcinoid syndrome patients present with initially normal 5-HIAA levels.⁴

The carcinoid syndrome may occur in the absence of liver metastases if the primary tumor originates from the bronchus^{17,18} or ovary^{11,19,20}; this is extremely rare in carcinoids of gastrointestinal origin.²¹ Although our patient did not display the intense prolonged flushing variant that has been described in some patients with bronchial carcinoids,²² a thorough evaluation did not reveal evidence of tumor other than the lung nodule before surgical intervention. Carcinoid tumors of the retroperitoneum and pancreas are rarely encountered in large series of carcinoid syndrome patients.^{1-4,23} To our knowledge, our patient's presentation with flushing due to an occult retroperitoneal carcinoid is unique.

The long-acting somatostatin analogue octreotide has received recent approval for marketing in the United States for the treatment of the carcinoid syndrome. Several studies have shown excellent response rates to octreotide in such patients, with reductions in flushing, diarrhea, and urinary 5-HIAA levels^{4,24,25} and a reversal of the carcinoid crisis.²⁶ Overt regression of carcinoid tumors with the use of octreotide has been reported but is relatively uncommon.^{4,24,25,27}

Our patient manifested a remarkable and sustained decrease in abdominal pain during octreotide therapy. Crampy abdominal pain in the carcinoid syndrome has been attributed to a direct bowel effect of serotonin²⁸; more intense pain is usually due to mesenteric retraction or tumor-related fibrosis with or without small bowel obstruction, mesenteric arterial insufficiency, or stretching of the liver capsule by metastatic tumor. Spontaneous necrosis of large liver metastases may present as an acute abdomen. This effect of octreotide therapy in our patient may relate to decreased hormoneogenesis or mesenteric tumor shrinkage below the limits of CT detection; alternatively, octreotide may modulate pain as a neurotransmitter in the peripheral or central nervous system.²⁹ Our results suggest that octreotide therapy should be considered in a patient with carcinoid with notable nonobstructive abdominal pain, even in the absence of troublesome flushing or diarrhea.

Steatorrhea appears to be a dose-related side effect of octreotide therapy^{24,30}; it rarely presents a clinical problem other than transiently bulky stools at the time of dosage escalations. Endogenous somatostatin excess is associated with gallstone formation,³¹ and gallstones and occasionally overt cholecystitis have been reported in patients treated long term

with octreotide.^{4,32,33} We recommend monitoring all patients during long-term octreotide therapy with periodic gall-bladder sonography.³³

A final clinical caveat is our patient's altered olfactory sensation at the onset of carcinoid symptoms with subsequent clear-cut subjective improvement after octreotide therapy. This may represent a new feature of the syndrome related to neurohumoral secretion³⁴ that has not been reported previously.

REFERENCES

1. Thorson AH: Studies on carcinoid disease. *Acta Med Scand* 1958; 161 (suppl 334):81-91
2. Grahame-Smith DG: *The Carcinoid Syndrome*. London, Heinemann Medical Books Ltd, 1972
3. Feldman JM: Carcinoid tumors and syndrome. *Semin Oncol* 1987; 14:237-246
4. Moertel CG: An odyssey in the land of small tumors. *J Clin Oncol* 1987; 5:1502-1522
5. Wilkin JK: Flushing reactions: Consequences and mechanisms. *Ann Intern Med* 1981; 95:468-476
6. Aldrich LB, Moattari AR, Vinik AI: Distinguishing features of idiopathic flushing and carcinoid syndrome. *Arch Intern Med* 1988; 148:2614-2618
7. Levine RJ, Sjoerdsma A: Pressor amines and the carcinoid flush. *Ann Intern Med* 1963; 58:818-828
8. Ahlman H, Dahlstrom A, Gronstad K, et al: The pentagastrin test in the diagnosis of the carcinoid syndrome: Blockade of gastrointestinal symptoms by ketanserin. *Ann Surg* 1985; 201:81-86
9. Norheim I, Theodorsson-Norheim E, Brodin E, et al: Tachykinins in carcinoid tumors: Their use as a tumor marker and possible role in the carcinoid flush. *J Clin Endocrinol Metab* 1986; 63:605-612
10. Pernow B, Waldenstrom J: Determination of 5-hydroxytryptamine, 5-hydroxyindole acetic acid and histamine in thirty-three cases of carcinoid tumor (argentaffinoma). *Am J Med* 1957; 34:16-25
11. Strodel WE, Vinik AI, Jaffe BM, et al: Substance P in the localization of a carcinoid tumor. *J Surg Oncol* 1984; 27:106-111
12. Feldman J, O'Dorisio TM: Role of neuropeptides and serotonin in the diagnosis of carcinoid tumors. *Am J Med* 1986; 81 (suppl 6B):41-48
13. Feldman JM: Urinary serotonin in the diagnosis of carcinoid tumors. *Clin Chem* 1986; 32:840-844
14. O'Connor DT, Deftos LJ: Secretion of chromogranin A by peptide-producing endocrine neoplasms. *N Engl J Med* 1986; 314:1145-1151
15. Feldman JM, Davis JA: Radioenzymatic assay of platelet serotonin, dopamine and norepinephrine in subjects with normal and increased serotonin production. *Clin Chim Acta* 1981; 109:275-283
16. Feldman JM: Serotonin metabolism in patients with carcinoid tumors: Incidence of 5-hydroxytryptophan-secreting tumors. *Gastroenterology* 1978; 75:1109-1116
17. VonBernheimer H, Ehringer H, Heistracher P, et al: Biologisch aktives, Nightmetastasierendes Bronchuscarcinoid mit linksherz Syndrom. *Wien Klin Wochenschr* 1960; 72:867-873
18. Todd TR, Cooper JD, Weissberg D, et al: Bronchial carcinoid tumors—Twenty years' experience. *J Thorac Cardiovasc Surg* 1980; 79:532-536
19. Sauer WG, Dearing WH, Flock EV, et al: Functioning carcinoid tumors. *Gastroenterology* 1958; 34:216-231
20. Kinley CE, Penner DW: Carcinoid tumors—A review and clinicopathologic study of 52 cases. *Can J Surg* 1962; 5:138-147
21. Feldman JM, Jones RS: Carcinoid syndrome from gastrointestinal carcinoids without liver metastasis. *Ann Surg* 1982; 196:33-37
22. Melmon KL, Sjoerdsma A, Mason DT: Distinctive clinical and therapeutic aspects of the syndrome associated with bronchial carcinoid tumors. *Am J Med* 1965; 39:568-581
23. Hodgson HJ, Maton PN: Carcinoid and neuroendocrine tumours of the liver. *Ballieres Clin Gastroenterol* 1987; 1:35-61
24. Kvols LK, Moertel CG, O'Connell MJ, et al: Treatment of the malignant carcinoid syndrome—Evaluation of a long-acting somatostatin analogue. *N Engl J Med* 1986; 315:663-666
25. Souquet JC, Sassolas G, Forichon J, et al: Clinical and hormonal effects of a long-acting somatostatin analogue in pancreatic endocrine tumors and in carcinoid syndrome. *Cancer* 1987; 59:1654-1660
26. Marsh HM, Martin JK Jr, Kvols LK, et al: Carcinoid crisis during anesthesia: Successful treatment with a somatostatin analogue. *Anesthesiology* 1987; 66:89-91
27. Wiedenmann B, R  th U, R  dsch R, et al: Tumor regression of an ileal carcinoid under the treatment with the somatostatin analogue SMS 201-995. *Klin Wochenschr* 1988; 66:75-77
28. Hendrix TR, Atkinson M, Clifton JA, et al: The effect of 5-hydroxytryptamine on intestinal motor function in man. *Am J Med* 1957; 23:886-893
29. Tolis G, Yotis A, del Pozo E, et al: Efficacy and safety of a somatostatin analogue in active acromegaly. In Reichlin S (Ed): *Somatostatin—Basic and Clinical Status*. New York, Plenum Press, 1987, pp 337-341
30. Lembcke B, Creutzfeldt W, Schleser S, et al: Effect of the somatostatin analogue sandostatin (SMS 201-995) on gastrointestinal, pancreatic, and biliary function and hormone release in normal man. *Digestion* 1987; 36:108-124
31. Krejs GJ, Orci L, Conlon JM, et al: Somatostatinoma syndrome: Biochemical, morphologic and clinical features. *N Engl J Med* 1979; 301:285-292
32. Comi RJ: Pharmacology and use in pituitary tumors. In Gordon P (Moderator): *Somatostatin and somatostatin analogue (SMS 201-995) in treatment of hormone-secreting tumors of the pituitary and gastrointestinal tract and non-neoplastic diseases of the gut*. *Ann Intern Med* 1989; 110:35-50
33. Jackson JA: Complications of long-acting somatostatin analog therapy (Letter). *Am J Med* 1989; 86:512-513
34. Jesberger JA, Richardson JS: Brain output dysregulation induced by olfactory bulbectomy: Approximation in the rat of major depressive disorder in humans? *Int J Neurosci* 1988; 38:241-265